

AD-A143 042

ON THE FILAMENTAL QUENCHING OF THE CURRENT-DRIVEN
ION-CYCLOTRON INSTABILITY (U) IOWA UNIV IOWA CITY DEPT
OF PHYSICS AND ASTRONOMY S L CARTIER ET AL JUN 84
U. OF IOWA-84-16 N00014-83-K-0452 F/G 20/7


1/1

UNCLASSIFIED

NL







MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A143 042

On the Filamental Quenching of the
Current-Driven Ion-Cyclotron
Instability

by

STEVEN L. CARTIER, NICOLA D'ANGELO,
PETER H. KRUMM,* and ROBERT L. MERLINO

Department of Physics and Astronomy
The University of Iowa
Iowa City, Iowa 52242

ABSTRACT

Experimental evidence is presented on the effect of the finite width of the current channel for the excitation of the current-driven ion-cyclotron instability. The results are in agreement with the non-local theory of Bakshi, Ganguli, and Palmadesso.

PACS numbers: 52.35.Fp, 52.35.Py

*Permanent address: Department of Physics, University of Natal,
Durban 4001, South Africa.

DTIC FILE COPY

SELECTED
JUL 13 1984
A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER U. of Iowa-84-16	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) On the Filamental Quenching of the Current-Driven Ion-Cyclotron Instability		5. TYPE OF REPORT & PERIOD COVERED Progress, June 1984
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Steven L. Cartier, Nicola D'Angelo, Peter H. Krumm, and Robert L. Merlino		8. CONTRACT OR GRANT NUMBER(s) N00014-83-K-0452
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Physics and Astronomy The University of Iowa Iowa City, Iowa 52242		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS --
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Physics Division Arlington, VA 22217		12. REPORT DATE June 1984
		13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Submitted to <u>Phys. Fluids</u>		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ion-Cyclotron Waves Nonlocal Theory Filamental Quenching		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Experimental evidence is presented on the effect of the finite width of the current channel for the excitation of the current-driven ion-cyclotron instability. The results are in agreement with the nonlocal theory of Bakshi, Ganguli, and Palmadesso.		

DD FORM 1473

1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Since their discovery over two decades ago,¹ ion-cyclotron waves have been an area of active research in the laboratory,² in the ionosphere-magnetosphere,³ and in the solar corona.⁴ The current-driven ion-cyclotron wave instability received its first theoretical treatment in the (local) theory of Drummond and Rosenbluth,⁵ appropriate to a uniform, magnetized plasma, without magnetic shear, in which electrons drift along \underline{B} field lines with the same drift velocity, \underline{v}_D , at all points in the plasma. Their treatment, however, could not allow, among other things, for the effect that the finite width of the current channel of most laboratory experiments has on the excitation of the instability. This has been considered in the nonlocal theories of, e.g., Ganguli and Bakshi⁶ and Bakshi, Ganguli, and Palmadesso.⁷ They find that if the width of the current channel is reduced to just a few ion Larmor radii, the instability is completely quenched. They refer to this phenomenon as filamental quenching.⁷ Indications of the presence of this phenomenon have been available since the first laboratory experiments on the ion-cyclotron wave instability and, more recently, from, e.g., the work of Sato.⁸

Here we report on a systematic test of the filamental quenching effect, performed on the single-ended Iowa Q-machine. A schematic diagram of the device is shown in Fig. 1(a). Plasma is produced by surface ionization of cesium atoms on a hot (~ 2200 K) tantalum plate 6 cm in diameter and is confined radially by a magnetic field of up to ~ 7 kG. The magnetic field is nonhomogeneous and varies along the



141

axis of the device into the end chamber, as shown in Fig. 1(b). The ion-cyclotron wave instability is excited, in the usual manner, by drawing current to a metallic disk moveable along the axis of the device, and is detected either in the current oscillations of the disk itself or by means of various (axially and radially moveable) Langmuir probes.

Figure 2 is a plot of the oscillation frequency versus disk position, as detected by the exciter disk or by a probe. The local ion-cyclotron frequency is also shown for comparison. The arrow indicates the location at which, for the particular conditions of Fig. 2, excitation of the fundamental mode ceases. By performing measurements similar to those of Fig. 2, but with various disk diameters and different currents in the magnet coils, we have been able to obtain the diagram shown in Fig. 3. Here the disk radius, r_D , is plotted as a function of r_i^* , the ion Larmor radius at the location of the disk, for which the instability is quenched. The line $r_D = r_i^*$ is also shown.

Evidently, the filamental quenching operates at widths of the current channel comparable to the local Larmor radius, in agreement with the conclusions of Bakshi, Ganguli, and Palmadesso⁶ (see, e.g., their Fig. 1). The small departure from the $r_D = r_i^*$ line for the 1.27 cm disk is most likely related to the large density perturbation introduced by this large disk.

Acknowledgments

We are grateful to Alfred Scheller for his enthusiastic technical support in the design and construction of the Iowa Q-machine. We also wish to thank G. Ganguli for useful discussions. This work was supported by a Northwest Area Foundation grant of the Research Corporation and by the United States Office of Naval Research under contract N00014-83-K-0452 and in part by NASA grant NGL-16-001-043.

References

- ¹N. D'Angelo and R. W. Motley, Phys. Fluids 5, 633 (1962); R. W. Motley and N. D'Angelo, Phys. Fluids 6, 296 (1963).
- ²See, for example, H. Böhmer, S. Fornaca, N. Rynn, and M. Wickham, Phys. Fluids 21, 2208 (1981); D. Dakin, T. Tajima, G. Benford, and N. Rynn, J. Plasma Phys. 15, 175 (1976); R. Schrittwieser, Phys. Fluids 26, 2250 (1983); R. Hatakeyama, N. Sato, H. Sugai, and Y. Hatta, Phys. Lett. 63A, 28 (1977).
- ³P. M. Kintner, M. C. Kelley, and F. S. Mozer, Geophys. Res. Lett. 5, 139 (1978); P. M. Kintner, M. C. Kelley, R. D. Sharp, A. G. Ghielmetti, M. Temerin, C. Cattell, P. F. Mizera, and J. F. Fennell, J. Geophys. Res. 84, 7201 (1979).
- ⁴See, for example, S. Hinata, Astrophys. J. 235, 258 (1980).
- ⁵W. E. Drummond and M. N. Rosenbluth, Phys. Fluids 5, 1507 (1962).
- ⁶G. Ganguli and P. Bakshi, Phys. Fluids 25, 1830 (1982).
- ⁷P. Bakshi, G. Ganguli, and P. Palmadesso, Phys. Fluids 26, 1808 (1983).
- ⁸N. Sato, Proceedings of the 1980 International Conference on Plasma Physics (Fusion Research Association of Japan, Nagoya, Japan, 1980), Vol. II, p. 138.

Figure Captions

Fig. 1. (a) Schematic cross-sectional view of the Iowa Q-machine.

(b) Axial variation of the B_z field in the region near and into the end chamber. For this case a coil current of 160 A produced a field $B_0 = 1170$ G in the center of the main chamber. The arrow indicates the $Z = 0$ position in parts (a) and (b). Electrostatic ion-cyclotron waves are produced by applying an appropriate bias to the axially moveable excitation disk, and are detected either in the disk current or by various Langmuir probes.

Fig. 2. Oscillation frequency versus exciter disk (0.64 cm radius) position (x). Oscillation frequency detected by a fixed Langmuir probe in the center of the main chamber, $Z = -47.5$ cm (\bullet). The arrow indicates the Z position at which excitation of the fundamental mode ceases. The solid line is the local ion-cyclotron frequency.

Fig. 3. Plot of exciter disk radius, r_D , versus r_1^* , the ion gyro-radius at the location of the disk for which the instability is quenched, for various magnet coil currents between 300 A and 800 A. The exciter disk radii used were: 0.32 cm, 0.64 cm, 0.79 cm, 0.95 cm, and 1.27 cm. The solid line is the line $r_D = r_1^*$.




Fig. 1




Fig. 2




Fig. 3

END

FILMED

8

1971







